

SEA SALT AEROSOL PRODUCTION: PARAMETERIZATION AND UNCERTAINTY

Ernie Lewis & Stephen E. Schwartz
elewis@bnl.gov ses@bnl.gov



Wednesday, December 15, 2004, 2:40 pm

American Geophysical Union
2004 Fall Meeting
San Francisco

IMPORTANCE of SEA SALT AEROSOL (SSA)

- Dominant aerosol species over the oceans
- One of the dominant aerosol species globally
- Light-scattering
 - Earth's radiation balance, global albedo
 - Remote sensing, visibility
- Properties of marine clouds
 - Anthropogenic radiative forcing – Twomey effect
 - Marine hydrology
- Atmospheric chemistry
 - Reactions within SSA particles
 - Nucleation suppression
- Air-sea exchange of latent heat, momentum, moisture

*Accurate parameterization of **size-dependent** SSA concentration and production flux (**including uncertainties**) is necessary.*

Sea Salt Aerosol Production


Mechanisms, Methods,
Measurements, and Models



Ernie R. Lewis and Stephen E. Schwartz

Available soon at your favorite bookseller (AGU)


SEA SALT AEROSOL PARTICLES AT 80% RH



r_{80} defined as radius of SSA particle in equilibrium at 80% RH

- r_{80} physically *relevant* (80% RH typical of marine boundary layer)
- r_{80} *unambiguous* (80% RH above hysteresis range)
- r_{80} *independent* of local meteorological conditions (RH)
- $r_{80} \approx 2r_{\text{dry}}$, $r_{80} \approx (1/2)r_{\text{form}}$


SEA SALT AEROSOL PARTICLES AT 80% RH




r_{80} defined as radius of SSA particle in equilibrium at 80% RH

- r_{80} physically *relevant* (80% RH typical of marine boundary layer)
- r_{80} *unambiguous* (80% RH above hysteresis range)
- r_{80} *independent* of local meteorological conditions (RH)
- $r_{80} \approx 2r_{\text{dry}}$, $r_{80} \approx (1/2)r_{\text{form}}$
- “Size” insufficient: $d_{\text{form}} = 2\{r_{\text{form}}\} \approx 2\{2[r_{80}]\} \approx 2\{[2(2r_{\text{dry}})]\}$


SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION




SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION



SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION




SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION




SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION

- *Shape* (size dependence of $n(r_{80})$) roughly independent of wind speed
- General *increase* in $n(r_{80})$ with increasing wind speed U_{10}
- Order of magnitude *variability* in $n(r_{80})$ at given U_{10}
- *Range of interest* $0.1 \mu\text{m} < r_{80} < 25 \mu\text{m}$
 - Few SSA particles with $r_{80} < 0.1 \mu\text{m}$
 - SSA particles with $r_{80} > 25 \mu\text{m}$ spend little time in the atmosphere
- Few data with $r_{80} < 1 \mu\text{m}$
- No evidence of multiple modes


SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION




SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION



SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION



SIZE DISTRIBUTION OF SSA NUMBER CONCENTRATION



PARAMETERIZATION of SIZE DISTRIBUTION of SSA CONCENTRATION

$$\left(\frac{dN}{d \log r_{80}} \right) / \text{cm}^{-3} = 0.07 \left(\frac{U_{10}}{\text{m s}^{-1}} \right)^2 \exp \left\{ -\frac{1}{2} \left[\frac{\ln(r_{80} / 0.3 \mu\text{m})}{\ln 2.8} \right]^2 \right\}$$

Associated uncertainty $\times 3$ (*essential*)


$$0.1 \mu\text{m} < r_{80} < 25 \mu\text{m}$$
$$5 \text{ m s}^{-1} < U_{10} < 20 \text{ m s}^{-1}$$

- Fits majority of data fairly well
- Lognormal - purely empirical
- Quadratic wind speed dependence; shape independent of wind speed
- Integral aerosol properties at $U_{10} = 10 \text{ m s}^{-1}$:
 - $N = (8 \times 3) \text{ cm}^{-3}$, mostly from $r_{80} = 0.1\text{-}1 \mu\text{m}$
 - $A = (80 \times 3) \mu\text{m}^2 \text{ cm}^{-3}$, mostly from $r_{80} = 1\text{-}10 \mu\text{m}$
 $\Rightarrow \sigma_{\text{sp}} \approx (40 \times 3) \text{ Mm}^{-1}$, $\tau_{\text{sp}} \approx 0.02 \times 3$
 - $M = (30 \times 3) \mu\text{g m}^{-3}$, mostly from $r_{80} = 3\text{-}20 \mu\text{m}$


PRINCIPAL METHODS of DETERMINING SSA PRODUCTION FLUX

<u>Method</u>	<u>Equation</u>	<u>r_{80} range</u>
Steady State Dry Deposition	$f_{\text{eff}}(r_{80}) = n(r_{80}) \cdot v_d(r_{80})$	$\sim 3 - \sim 25 \mu\text{m}$
Statistical Wet Deposition	$f_{\text{eff}}(r_{80}) = n(r_{80}) \cdot H_{\text{mbl}} / \tau_{\text{wet}}$	$< \sim 1 \mu\text{m}$
Concentration Buildup	$f_{\text{eff}}(r_{80}) = U \frac{\partial}{\partial x} \left[\int n(r_{80}, x, z) dz \right]$	$< \sim 10 \mu\text{m}$
Micrometeorological (Eddy correlation)	$f_{\text{eff}}(r_{80}) = \overline{n'(r_{80}) w'}$	$< \sim 1 \mu\text{m}$
Whitecap	$f_{\text{int}}(r_{80}) = f_{\text{wc}}(r_{80}) \cdot W(U_{10})$	$< \sim 10 \mu\text{m}$


EFFECTIVE SSA PRODUCTION FLUX




EFFECTIVE SSA PRODUCTION FLUX




EFFECTIVE SSA PRODUCTION FLUX




EFFECTIVE SSA PRODUCTION FLUX




EFFECTIVE SSA PRODUCTION FLUX




EFFECTIVE SSA PRODUCTION FLUX




EFFECTIVE SSA PRODUCTION FLUX



EFFECTIVE SSA PRODUCTION FLUX



EFFECTIVE SSA PRODUCTION FLUX



PARAMETERIZATION of SIZE DISTRIBUTION of EFFECTIVE SSA PRODUCTION FLUX


$$\left(\frac{dF_{\text{eff}}}{d \log r_{80}} \right) / \left(\text{m}^{-2} \text{s}^{-1} \right) = 50 \left(\frac{U_{10}}{\text{m s}^{-1}} \right)^{2.5} \exp \left\{ -\frac{1}{2} \left[\frac{\ln (r_{80} / 0.3 \mu\text{m})}{\ln 4} \right]^2 \right\}$$

Associated uncertainty $\times 5$ (*essential*)

$$0.1 \mu\text{m} < r_{80} < 25 \mu\text{m}$$
$$5 \text{ m s}^{-1} < U_{10} < 20 \text{ m s}^{-1}$$

- Purely empirical
- Lognormal shape (i.e., size dependence) independent of wind speed

SUMMARY



Size $r_{80} \approx 2r_{\text{dry}} \approx (1/2)r_{\text{form}}$

- Range of interest $r_{80} \approx 0.1\text{-}25 \mu\text{m}$

SSA concentration $n(r_{80})$ – uncertainty for given U_{10} is $\times 3$

- Concentrations increase roughly as U_{10}^2
- Shape of size distribution roughly independent of wind speed
 - **Number** concentration mostly $r_{80} \approx 0.1\text{-}1 \mu\text{m}$ (CCN)
 - **Area** concentration mostly $r_{80} \approx 1\text{-}10 \mu\text{m}$ (light scattering)
 - **Mass** concentration mostly $r_{80} \approx 3\text{-}20 \mu\text{m}$

Effective SSA production flux $f_{\text{eff}}(r_{80})$ – uncertainty for given U_{10} is $\times 5$

- Flux increases roughly as $U_{10}^{2.5}$
- Shape of size distribution roughly independent of wind speed

<http://www.ecd.bnl.gov/steve/SSA-AGUfm04.pdf>

Sea Salt Aerosol Production

Mechanisms, Methods,
Measurements, and Models



Ernie R. Lewis and Stephen E. Schwartz

Another shameless plug for our book. Order now at the AGU booth!